

International Journal of Medical Scholars

Article DOI:

Volume 1, Issue 1, March 2023



Horizontal mattress suture for intestinal anastomosis

Publication History

Received: Jan 01, 2023 Revised: Jan 11, 2023

Accepted: Feb 04, 2023 Published: March 30, 2023

Authors and Affiliation:

Attaullah Mahar^{1*}, Saim Ather², Muhammad Osama Rauf Hiraj³, Muhammad Sabir⁴, Saifullah Brohi⁵

¹National Medical Center, Karachi, Pakistan

² Ibn-e-Sina Hospital, Multan, Pakistan

^{3,4}Nishtar Hospital, Multan, Pakistan

⁵Bilawal Medical College for Boys at LUMHS
Jamshoro, Pakistan

*Corresponding Author Email:

drattamahar2@gmail.com

Copyright & Licensing:



Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a [Creative Commons Attribution \(CC-BY\) 4.0 License](https://creativecommons.org/licenses/by/4.0/) that allows others to share the work with an acknowledgment of the work's authorship and initial publication in this journal.

Conflict of Interest:

Author(s) declared no conflict of interest.

Acknowledgment:

No Funding received.

Citation: Mahar A, Ather S, Hiraj MOR, Sabir M, Brohi S. Horizontal mattress suture for intestinal anastomosis. International Journal of Medical Scholars. 2023 March 30; 1(1):1-6.

Please scan me to access online.



An official publication of
Medteach Private Limited, Multan, Pakistan.

Email: info@medteach.org, Website: <https://www.medteach.org/index.php/ijms>



International Journal of Medical Scholars

Volume 1, Issue 1, 2023; pp: 1-6

Original Article



Horizontal mattress suture for intestinal anastomosis

Attaullah Mahar^{1*}, Saim Ather², Muhammad Osama Rauf Hiraj³, Muhammad Sabir⁴, Saifullah Brohi⁵

¹National Medical Center, Karachi, Pakistan

² Ibn-e-Sina Hospital, Multan, Pakistan

^{3,4}Nishtar Hospital, Multan, Pakistan

⁵Bilawal Medical College for Boys at LUMHS Jamshoro, Pakistan

*Corresponding Author Email: drattamahar2@gmail.com

ABSTRACT

Objective: Investigating the use of a new horizontal mattress suture anastomosis technique for patients who have undergone emergency or elective intestinal anastomosis.

Methods: In this prospective study a total of 200 patients were included in the study, 100 managed with horizontal mattress suturing technique and 100 were treated with traditional double layer method. Main variables of study were anastomosis time, hospital stay, CRP, post-operative infection, anastomosis leakage, reintervention and readmission in hospital.

Results: The mean anastomosis time of traditional 2 layer patients was greater than horizontal mattress suture patients as 19.68 ± 2.01 minutes and 13.67 ± 2.38 minutes, respectively. ($p < 0.010$). Regular diet tolerated within POD 4, CRP at POD 5 and postoperative hospital stay in both the groups were almost equal, ($p > 0.010$).

Conclusion: Horizontal mattress suture is simpler and associated with fewer complications, it may lead to shorter postoperative hospital stays in patients who were planned for intestinal anastomosis/laparotomy either emergency or elective. This could be beneficial for both patients and healthcare providers by reducing costs and the risk of hospital-acquired infections.

Keywords: Horizontal mattress anastomosis, Abscess, Anastomotic leakage, Two-layer anastomosis, Intestinal repair

1. INTRODUCTION

Anastomosis is a surgical procedure where two ends of a tubular structure, such as the digestive tract, are connected or joined together¹. Traditional manual anastomosis and surgical staples are two common methods used for this purpose, but they each have their own advantages and disadvantages². Traditional manual technique has advantage of precise control, low risk of foreign body reaction and disadvantage of potential for leakage³. On other hand stapled technique has advantage of fast procedure with consistency and handling with less experience and disadvantages include intestinal fistula, anastomosis leak and limited customization⁴. Compression of soft tissue when using traditional stitching methods, is indeed a valid concern in certain medical and surgical contexts⁵. When sutures are applied with excessive tension or pressure, they can compress and damage the surrounding soft tissues⁶. Partial incisions and tearing during a surgical procedure, particularly in the context of an anastomosis can have serious consequences like leakage, bleeding, impaired healing and infection risk⁷.

Another renowned procedure is horizontal mattress suture technique commonly used in various surgical procedures, including intestinal reconstruction⁸. This technique is particularly useful when there is a need to securely approximate tissues with good tension distribution and knot security⁹. In intestinal surgery, it can be used to close intestinal incisions or anastomoses. The advantages of the mentioned approach over other techniques include simplicity; ease of execution, reduced operating time, wide applicability and it is particularly suited for vessel anastomosis in ischemic or edematous tissues¹⁰.

We compared two different methods of gastrointestinal anastomosis in adult patients the whole wall horizontal mattress suture anastomosis and the traditional manual anastomosis method. The aim of the study was to investigate the efficacy of this novel manual method in gastrointestinal anastomosis.

2. METHODOLOGY

Study was conducted at department of general surgery, Nishtar hospital, Multan, from 31st July 2022 to 30th June 2023 in one year duration. Study was approved by hospital ethical board after complete investigation and clarification of study purpose. Non probability consecutive sampling technique was used and sample size was calculated by using online sample size calculator with 95% confidence interval, 80% study power and readmission in traditional method was 55% and 46% in horizontal suture method. Patients with liver injury diagnosed intra-operatively or by CT scan, both gender and age 16-60 years were included in the study.

Patients who require primary bowel resection and intestinal anastomosis, age 16-60 years, both genders, normal renal function, no sepsis and administration of immunosuppressive and steroid drugs were included. Patients having anastomosis of duodenum, rectum and stomach. Patients managed in intensive care in more than 3 days. Bowel preparation, antibiotic prophylaxis, and a suture technique using a monofilament suture material was done under general anesthesia. This part of the passage refers to the technique used to reconnect the remaining sections of the bowel after the diseased segment has been removed. "End-to-end" and "end-to-side" refer to different methods of creating the new connection or anastomosis between the two ends of the bowel.

The posterior wall was constructed using a continued horizontal mattress anastomosis, and the anterior wall was sutured with Connell sutures. These techniques are used to ensure a secure and effective closure of tissue layers during surgery. An edge distance of approximately 2 to 3mm means that the needle is inserted about 2 to 3mm away from the edge of the tissue. During surgical connection of two blood vessels or tubular structures (e.g., intestines), ischemia must be avoided by putting low pressure to ensure that the connected tissues receive sufficient blood supply to remain healthy.

Every patient who underwent a surgical procedure followed the same set of postoperative care guidelines or steps including intravenous fluid, nutrition support, antibiotics administration and routine follow up. Collected data was analyzed with SPSS version 24. Test of significance was applied and p value ≤ 0.05 was taken as significant.

3. RESULTS

Overall, 200 patients were included in our study both genders. The patients were equally divided into two groups as horizontal mattress suture 100 (50.0%) and traditional 2 layer 100 (50.0%). The mean age of horizontal mattress suture and traditional 2 layer patients was 45.36 ± 12.69 years and 46.91 ± 11.19 years, respectively. The most common cause of operation in both the groups was intussusception, 34 (34.0%) and 42 (42.0%), respectively. The differences were statistically insignificant, ($p > 0.010$). (Table. I).

The mean anastomosis time of traditional 2 layer patients was greater than horizontal mattress suture patients as 19.68 ± 2.01 minutes and 13.67 ± 2.38 minutes, respectively. ($p < 0.010$). Regular diet tolerated within POD 4, CRP at POD 5 and

postoperative hospital stay in both the groups were almost equal, ($p > 0.010$). (Table. II).

Postoperative complications occurred in both the groups were presented in figure. I.


Table-I: Demographic and baseline of the study groups

Variable	Propensity matched patients		p-value
	Horizontal mattress suture	Traditional 2 layer	
Age	45.36±12.69	46.91±11.19	0.755
Gender			
Male	71 (71.0)	66 (66.0)	0.447
Female	29 (29.0)	34 (34.0)	
Causes of operation			
Intussusception	34 (34.0)	42 (42.0)	0.058
Trauma	29 (29.0)	38 (38.0)	
Perforation	19 (19.0)	12 (12.0)	
Obstruction/strangulation	18 (18.0)	8 (8.0)	

Table. II: Surgical outcome of the study groups

Variable	Propensity matched patients		p-value
	Horizontal mattress suture	Traditional 2 layer	
Anastomosis time (min)	13.67±2.38	19.68±2.01	<0.010
Regular diet tolerated within POD 4	22 (22.0)	28 (28.0)	0.327
CRP at POD 5 (mg/L)	16.21±3.41	16.83±3.21	0.186
Postoperative hospital stay (days)	10.01±2.23	11.38±2.11	0.331

Figure. I



4. DISCUSSION

The ability of an intestinal anastomosis technique to heal without leakage is indeed the ultimate test of its suitability. Leakage at the anastomotic site can lead to a range of serious complications including, sepsis, peritonitis, delayed healing, abscess formation, nutritional issues and reoperation¹¹. In our study surgical wound infection is common complication found in 23% of patients 20% in horizontal sutures and 23% in traditional suture technique. Second common complication was hospital readmission which was also higher in traditional suture method.

Placer et al¹² conducted a study and concluded that anastomotic leaks present with immediate, dramatic symptoms. Some patients may experience a more insidious or gradual onset of symptoms (low grade fever, failure to thrive and prolonged ileus). In another study Hyman et al¹³ reported that postoperative abscess can indeed be caused by a small anastomotic leak. Various reasons, such as infection, poor surgical technique, or inadequate postoperative care are responsible for it.

Yang et al¹⁴ conducted a study on pediatric population and reported that manual anastomosis technique has shown beneficial effects in pediatric patients in terms of reduced anastomotic complications and shorter hospital stays. Pediatric patients and their families may experience improved quality of life due to reduced hospitalization. Basar et al¹⁵ also concluded that continuous horizontal mattress suture technique may reduce the time required for anastomosis compared to other suturing techniques. The horizontal mattress pattern allows for quicker and more efficient suturing in some cases.

By eversion of the tissue and careful placement of sutures, the continuous horizontal mattress suture helps create a smoother, more uniform inner surface

(intimal surface) at the anastomotic site¹⁶. This approach contributes to better patient outcomes in surgeries involving tubular structures like blood vessels and the gastrointestinal tract. This suturing technique with its everting effect provides more perfect microsurgical anastomosis which is an addition to other advantages of this procedure¹⁷.

Suturing is a fundamental skill in medicine and surgery, and the choice of sutures and suturing techniques can significantly impact the healing process. Horizontal mattress suturing is also promotes healing process and reduced complications proved in dental surgeries¹⁸. Controversial study findings were also reported by Radad et al¹⁹ who preferred simple interrupted suture technique because of its patency rate. In another study Chung et al²⁰ reported that the horizontal mattress running stitches with a single monofilament suture, may indeed have been considered safe for certain medical procedures, such as wound closure or surgical suturing. The safety and effectiveness of a specific technique can depend on various factors, including the type of procedure, the patient's individual circumstances, and the surgeon's skill.

5. CONCLUSION

Horizontal mattress suture is simpler and associated with fewer complications, it may lead to shorter postoperative hospital stays in patients who were planned for intestinal anastomosis/laparotomy either emergency or elective. This could be beneficial for both patients and healthcare providers by reducing costs and the risk of hospital-acquired infections.

Limitations: Performing propensity score matching analysis is a valuable statistical technique in research and epidemiology to reduce the impact of potential confounding variables when studying the effects of a

particular intervention or treatment, such as a modified anastomotic method.

6. REFERENCES

1. Leahy J, Schoetz D, Marcello P, Read T, Hall J, Roberts P, et al. What is the risk of clinical anastomotic leak in the diverted colorectal anastomosis? *J Gastrointest Surg.* 2014;18:1812–6.
2. Maemura K, Mataka Y, Kurahara H, Tanoue K, Kawasaki Y, Ijichi T, et al. The 1-year outcomes after pancreaticogastrostomy using vertical versus horizontal mattress suturing for gastric wrapping. *Surg Today.* 2021;51:511-9.
3. Millano L, Fadhila I, Anita M. Primary esophageal anastomotic using mattress suturing technique in cases of esophageal atresia with fistula. *Intern J Surg.* 2023;7(1):88-91.
4. Ghosh A, Biswas SK, Ghosh T, Basu KS, Biswas SK. Pediatric colonic anastomosis: Can method of anastomosis and wound closure be of help?. *J Indian Assoc Pediatr Surg.* 2023;28(3):245.
5. Sumner SM, Regier PJ, Case JB, Ellison GW. Evaluation of suture reinforcement for stapled intestinal anastomoses: 77 dogs (2008-2018). *Veterinary Surg.* 2019;48(7):1188-93.
6. Hobbs RD, Moon J, Murala J, Ohye RG. Novel suture technique for slide tracheoplasty for the treatment of long-segment tracheal stenosis. *Semin Thorac Cardiovasc Surg.* 2020 Winter;32(4):930-934.
7. Takagi K, Domagala P, Polak WG, Buettner S, Ijzermans JNM. The controlling nutritional status score and postoperative complication risk in gastrointestinal and hepatopancreatobiliary surgical oncology: a systematic review and meta-analysis. *Ann Nutr Metab.* 2019;74:303–12.
8. Sliker JC, Daams F, Mulder IM, et al. Systematic review of the technique of colorectal anastomosis. *JAMA Surg.* 2013;148:190–201.
9. Byrne M, Aly A. The Surgical Suture. *Aesthet Surg J.* 2019 Mar 14;39(Suppl_2):S67-S72.
10. Feng SM, Han M, Wang AG, Fan JQ. Functional comparison of horizontal mattress suture versus free-edge suture in the all-inside arthroscopic Broström–Gould procedure for chronic lateral ankle instability. *Orthop Surg.* 2020;12(6):1799-1810.
11. Harris LJ, Moudgill N, Hager E, Abdollahi H, Goldstein S. Incidence of anastomotic leak in patients undergoing elective colon resection without mechanical bowel preparation: our updated experience and two-year review. *Am Surg.* 2009;75:828–33.
12. Placer C, Enríquez-Navascués JM, Elorza G, Timoteo A, Mugica JA, Borda N, et al. Preventing complications in colorectal anastomosis: results of a randomized controlled trial using bioabsorbable staple line reinforcement for circular stapler. *Dis Colon Rectum.* 2014;57:1195–1201.
13. Hyman NH, Osler T, Cataldo P, Burns EH, Shackford SR. Anastomotic leaks after bowel resection: what does peer review teach us about the relationship to postoperative mortality? *J Am Coll Surg.* 2009; 208:48–52.
14. Yang Y, Chen B, Xiang L, Guo C. Reduced rate of dehiscence after implementation of a novel technique for creating colonic anastomosis in

- pediatric patients undergoing intestinal anastomosis in a single institute. *Medicine (Baltimore)*. 2019 May;98(19):e15577.
15. Başar H, Erol B, Tetik C. Use of continuous horizontal mattress suture technique in end-to-side microsurgical anastomosis. *Hand Surg*. 2012;17(03):419-27.
 16. QU Y, MAN Y. Key points to execute internal horizontal mattress suture in tissue augmentation. *J Prevention Treatment Stomatol Dis*. 2023:153-61.
 17. Connaughton AJ, Kluczynski MA, Marzo JM. Simple versus horizontal mattress suture configuration in bankart repair. *J Orthop*. 2021;23:225-6.
 18. Mahabaleshwara CH, Francis M, CR TA, Pethkar R, Shravani B. Simple interrupted and continuous interlocking suturing techniques comparison in maxillofacial trauma—a randomized controlled trial. *RGUHS J Dental Sci*. 2021;13(3):198-201.
 19. Radad K, El-Shazly M. Clinical and pathological assessment of different suture techniques for microvascular anastomosis in rat femoral artery. *J Veterinary Sci*. 2007;8(3):269-73.
 20. Chung RS, Hitch DC, Armstrong DN. The role of tissue ischemia in the pathogenesis of anastomotic stricture. *Surg*. 1988;104:824–9.